

# Antimicrobial Activity of Medicinal Plants: A Weapon to Combat Multi-Drug Resistant Pathogens

Mobina Parveen<sup>1</sup>, Sudip Some<sup>2\*</sup>

DOI: 10.18811/ijpen.v7i02.3

## ABSTRACT

Medicinal plants are extensively used in traditional systems of medicine for the treatment of different ailments. Bio-active compounds of plants and herbs show excellent antimicrobial properties against different pathogens. Plant extracts are known to control microbial growth by their anti-quorum sensing and anti-biofilm formation activities. The emergence and rapid spread of antibiotic resistance in bacterial pathogens and the evolution of new multi-drug resistant strains of disease-causing microorganisms are of remarkable alarm to the healthcare systems across the globe. Therefore the antimicrobial properties of the plant extract may represent an important alternative therapeutic route to conventional drugs. Phytochemical analysis of extract also exclusively facilitates understanding their biochemical and molecular nature for drug discovery shortly. The article focuses on the antimicrobial activity of some selected medicinal plants against the most common microbial pathogens, including clinical multi-drug resistant (MDR) bacteria.

**Keywords:** Antimicrobial activity, Bio-active compounds, MDR bacteria, Medicinal plants, Traditional medicine.

*International Journal of Plant and Environment* (2021);

ISSN: 2454-1117 (Print), 2455-202X (Online)

## INTRODUCTION

Since the dawn of civilization man and plants have had a deep relationship for food, timber, spices, and drugs. Ethnic communities across the globe have been enriched by their own culture in the form of traditional healthcare systems. Indian traditional systems of medicine are based on Ayurveda, Yoga, Unani, Siddha, and Homoeopathy (Prasad, 2002; Ravishankar and Shukla, 2007; Neba, 2011; Teka *et al.*, 2020). Different physical ailments and their treatments have been cited in the great *Charaka Samhita*, the ancient Indian comprehensive medical text of Ayurveda (Bhavana and Shreevathsa, 2014). According to World Health Organization (WHO), about 75% of the world's population and about 80% population of the developing countries depend directly on herbal drugs for their primary healthcare. Different parts of the medicinal plants, including roots, leaves, fruits, seeds, or whole plants, have been used as therapeutic herbal drugs. Traditional healers or *Baidhya* are advising to take the folk medicines through different ethno-medicinal formulations such as plant extract, decoction, paste, powder, and infusion for its excellent anti-inflammatory, anti-diarrhoeal, antioxidant, and antimicrobial properties (Uporey *et al.*, 2012; Dolatkahi *et al.*, 2014; Pan *et al.*, 2014; Some *et al.*, 2017). The traditional systems of medicine are getting advantages in healthcare due to fewer side effects, excellent curative treatment for several chronic diseases, and high cost of new drugs (Pandey *et al.*, 2013; Some and Mukherjee, 2018). Most of the plants contain a variety of bio-active compounds. Ethno-botany and ethno-medicinal investigations are now acknowledged most efficient techniques for updating or refocusing the data on medicinal plants reported in previous research for the possible extraction of useful bio-active compounds (Thirumalai *et al.*, 2009). These phytochemicals showed excellent antimicrobial activity against various Gram-positive and Gram-negative bacteria (Mahasneh and El-Oqlah, 1999; Srinivasan *et al.*, 2001; Das *et al.*, 2010). In addition to the phytochemicals in plant extract inhibits microbial growth by interfering cell-cell communication processes, which is generally

<sup>1</sup>Department of Botany, Dhruba Chand Halder College, Dakshin Barasat, South 24-Paraganas-743372, West Bengal, India

<sup>2</sup>Department of Life Sciences, Chanchal Siddheswari Institution (H.S.), Chanchal, Malda-732123, West Bengal, India

**\*Corresponding author:** Sudip Some, Department of Life Sciences, Chanchal Siddheswari Institution (H.S.), Chanchal, Malda-732123, West Bengal, India, Mobile: +91-9474347483, Email: ssome.csi@gmail.com

**How to cite this article:** Parveen, M. & Some, S. (2021). Antimicrobial Activity of Medicinal Plants: A Weapon to Combat Multi-Drug Resistant Pathogens. *International Journal of Plant and Environment*, 7(2), 133-142.

**Conflict of interest:** None

**Submitted:** 23/05/2021 **Accepted:** 30/06/2021 **Published:** 15/07/2021

termed as Quorum Sensing (QS) (Bacha *et al.*, 2016). Several plant antimicrobial peptides (AMPs) in roots, seeds, flowers, stems, and leaves showed wonderful antimicrobial effect against different viruses, bacteria, fungi, protozoa, parasites by interaction with phospholipids and membrane permeabilization (Nawrot *et al.*, 2014). The emergence of antibiotic resistance in microorganisms establishes a serious health issue due to several factors, including misuse and overuse of antibiotics. The association of antibiotic resistance genes with many mobile genetic components including plasmids, transposons, integrons, and gene cassettes, has facilitated the rise of drug resistance in bacteria. Steps can be taken at all levels of society to reduce the impact and limit the spread of resistance (White, 2009; Elisha *et al.*, 2017).

Therefore, the scientific community has focused their keen interest in pharmacognosy where different bioactive compounds are being screened from plant extract to use as an alternative medicine to combat microbial infection. In this context, various researches have been reported simultaneously in the literature. Full elaboration goes beyond the reach of this inquiry as the present review article is unique to selected plants or microbes. The present article focuses on traditional uses of medicinal plants, bioactive compounds in medicinal plants, and the promising antimicrobial activity of plant extract.

## TRADITIONAL USES OF MEDICINAL PLANTS

Ethno-botanical research is indispensable to exploring the traditional uses of plants for medicinal purposes. In this context, Tefera and Kim (2019) explored the traditional uses of medicinal plants among the ethnic communities in Southern Ethiopia. They observed that 105 medicinal plants belonging to 52 families and 96 genera had been used to treat human and livestock ailments, including gastrointestinal, dermatological, skeletomuscular, respiratory, and parasitic diseases. Most of the plants of Fabaceae family followed by Lamiaceae family have been utilized in primary healthcare system. Another ethno-botanical survey was conducted for the documentation of traditional knowledge in Northeastern Ethiopia. A total of 91 medicinal plant species belonging to 51 families have been used to cure 38 human diseases and 12 livestock ailments. The leaf is the most frequently used plant part in the preparation

of herbal drug. *Ocimum urticifolium* Roth has been used as the most preferred plant species in therapeutic practice followed by *Withania somnifera* (L.) Dunal and *Zehneria scabra* Sond. (Osman *et al.*, 2020). An ethno-medicinal study has shown that *Euclea divinorum* Hiern, *Tylosema fassoglensis* (Schweinf.) Torre & Hillc., *Carissa edulis* (Forssk.) Vahl, *Harrisonia abyssinica* Oliv., *Zanthoxylum gillettii* (De Wild.) P.G. Waterman, and *Warburgia salutaris* (G. Bertol.) Chiov. have been predominantly used in respiratory diseases by the people of Kisumu East Sub County of Kenya (Mailu *et al.* 2020). Some popular therapeutic plants that have been used traditionally are presented in Table 1.

## BIO-ACTIVE COMPOUNDS IN MEDICINAL PLANTS

Medicinal plants produce bio-active compounds due to secondary cellular metabolism classified into four major

**Table 1:** Traditional uses of medicinal plants.

Botanical Name	Family	Use Parts	Medicinal Uses	References
<i>Aloe vera</i> L.	Asphodelaceae	L	Leaf extract has been used in dermal problems.	Abdelhalim <i>et al.</i> , 2017
<i>Amaranthus graecizans</i> L.	Amaranthaceae	L	Leaves have been used as laxative.	Mahmoud and Gairola, 2013
<i>Boerhavia diffusa</i> L.	Nyctaginaceae	L, R	Extract of leaves and roots have been used in gastrointestinal, respiratory, cardiovascular, and nephrological problems.	Mishra <i>et al.</i> , 2014
<i>Bryophyllum pinnatum</i> (Lam.) Kurz.	Crassulaceae	L	Leaf extract has been used to cure wounds and burns.	Saini <i>et al.</i> , 2016
<i>Caesalpinia decapetala</i> (Roth.) Alston	Fabaceae	R	Decoction of root has been used in menstrual problems.	Mahwasane <i>et al.</i> , 2013
<i>Cannabis sativa</i> L.	Cannabaceae	S	Seed extract has been applied in rheumatoid problem, neurological, and rectal problems.	Uniyal <i>et al.</i> , 2006
<i>Chenopodium album</i> L.	Chenopodiaceae	Wp	Plant extract has been used in gastrointestinal and nephrological problems.	Aziz <i>et al.</i> , 2018
<i>Eucalyptus obliqua</i> L'Hér.	Myrtaceae	L	Decoction of leaf has been applied to cure upper respiratory infection.	Morales <i>et al.</i> , 2016
<i>Justicia adhatoda</i> L.	Acanthaceae	L	Leaf extract has been used in respiratory problems.	Vidarthi <i>et al.</i> , 2013
<i>Mentha piperita</i> L. Huds.	Lamiaceae	L	Decoction of leaves has been used in gastrointestinal troubles.	Abdelhalim <i>et al.</i> , 2017
<i>Mimosa pudica</i> L.	Mimosaceae	R	Root extract has been used in urinary incontinence and bladder infection.	Nandagoapalan <i>et al.</i> , 2015
<i>Momordica charantia</i> L.	Cucurbitaceae	F	Fruit extract has been used as anti-diabetic and blood purifier.	Mahomoodally, 2013
<i>Smilax aspera</i> Wall.	Smilacaceae	R	Root extract has been used in sexually transmitted diseases and dermal problems.	Vidarthi <i>et al.</i> , 2013
<i>Solanum indicum</i> Burm.	Solanaceae	L, R	Root is dried to form powder and blended to form paste with lemon juice. It has been used in eye disease. Leaf extract has been used for cold and cough.	Vidarthi <i>et al.</i> , 2013
<i>Vitex negundo</i> L.	Verbenaceae	L	Leaves are boiled in water. Steam is inhaled for cold, cough, and sinusitis.	Vidarthi <i>et al.</i> , 2013


Abbreviations: F: Fruit; L: Leaf; R: Root; S: Stem; Wp: Whole plant

classes like alkaloids, glycosides, polyphenols, and terpenoids (Ahmed *et al.*, 2017). The foremost phytochemicals isolated from *Chenopodium album* L. include phenols, alkaloids, glycosides, saponins, and flavonoids, having an excellent pharmacological activities (Poonia, 2021). The primary phytochemical screening of the leaf extract of *M. pudica* L. exhibited the existence of bio-active components such as terpenoids, flavonoids, glycosides, alkaloids, quinines, phenols, tannins, saponins, and coumarins (Gandhiraja *et al.*, 2009; Rajendran and Sundararajan, 2010; Ahmad *et al.*, 2012). Apart from antimicrobial activity, flavan-3-ols from cocoa (*Theobroma cacao* L.) plant are related to a reduced risk of cardio-vascular diseases, diabetes mellitus, insulin resistance, as well as improvements in lipids, endothelial-dependent blood flow and blood pressure and systemic inflammation (Fraga *et al.*, 2019). It has been noted that the yield of non-polar phytochemicals in aqueous extract was recorded quite lower than the alcoholic extract due to polarity differences between the two solvents. The plant extract of *Cynodon dactylon* (L.) Pers exhibited 12.2% glycosides, 6.3% tannins, 0.1% alkaloids, 1.0% resins, 10% free reducing sugar, and 12% total reducing sugar, respectively (Jolly and Narayanan, 2000). A study has shown that the values of terpenoids & phenolics, alkaloids, quaternary alkaloids & N-oxides and fats & waxes were recorded as 5.54067%, 2.3673%, 17.3223%, and 1.6733%, respectively in the plant extract of *Tridax procumbens* (L.) L. (Shaikh, 2018). In another study, Fahal *et al.* (2018) measured the amounts of alkaloids, flavonoids, saponins, sterol, and tannins with the values of 3.1% ( $\pm 0.004$ ), 5.2 % ( $\pm 0.003$ ), 6.5%

( $\pm 0.1$ ), 4.7% ( $\pm 0.002$ ), and 22.9% ( $\pm 0.01$ ), respectively in the crude extract of the pod of *Moringa oleifera* Lam. The name and chemical structures of some common bio-active compounds are presented in Fig 1. The qualitative and quantitative data of bio-synthesized active compounds in medicinal plants are displayed in Table 2. Several researchers have screened the phytochemicals from different medicinal plants using various analytical techniques. A brief account of phytochemicals in medicinal plants is presented in Table 3.

## ANTIMICROBIAL ACTIVITY

The secondary metabolites or bioactive compounds in medicinal plants have excellent therapeutic potency against microbes. Tannins, terpenoids, alkaloids, flavonoids, and saponins exhibited excellent antimicrobial activity against bacteria, fungi, and protozoa (Oladeji, 2016). The protocol of plant extract-mediated antimicrobial activity is represented in Scheme 1.



**Fig 1:** Chemical structure of bioactive compounds in the medicinal plant (The figure has been adapted from Mngeni, 2017).


**Table 2:** Qualitative and quantitative data of bio-synthesized active compounds in medicinal plants.

Name of medicinal plants	Bio-active compounds	Quantity (mg of GAE or QE /g of dry weight)	References
<i>Aegle marmelos</i> (L.) Corrêa.			
<i>Cynodon dactylon</i> (L.) Pers.			
<i>Eclipta prostrata</i> (L.) L.	Alkaloids, flavonoids, saponins, steroid, tannins, terpenoid, phlobatannin, and cardiac glycoside.	-	Dhandapani and Sabna, 2008
<i>Moringa pterygosperma</i> Gaertn.			
<i>Pongamia pinnata</i> (L.) Pierre			
<i>Sida acuta</i> Burm. f.			
<i>Tridax procumbens</i> (L.) L.			
<i>Aegle marmelos</i> (L.) Corrêa	Flavonoids	166.33 ( $\pm 09.6$ )	Rajan <i>et al.</i> , 2011
	Tannins	193.33 ( $\pm 22.03$ )	
	Phenolic compounds	158.66 ( $\pm 28.67$ )	
<i>Sida acuta</i> Burm. F	Total flavonoids in leaf extract	8.5	Jindal <i>et al.</i> , 2012
	Bound flavonoids in bud extract	1.75	
<i>Eclipta alba</i> (L.) Hassk	Total phenolics	31.67 ( $\pm 3.82$ )	Yadav <i>et al.</i> , 2017
	Flavonoids	91.67 ( $\pm 5.05$ )	
<i>Pongamia pinnata</i> (L.)	Total flavonoids	0.01398	Mishra and Pounikar, 2019
<i>Rhodomertus tomentosa</i> (Aiton) Hassk.	Total phenolics	0.7377	
<i>Ziziphus lotus</i> (L.) Lam.	Total phenolics	9.4987	Yahia <i>et al.</i> , 2020
	Total flavonoids	0.92055	
<i>Ziziphus mauritiana</i> Lam.	Total phenolics	5.3295	
	Total flavonoids	0.9026	

[GAE: Gallic Acid Equivalent; QE: Quercetin Equivalent]

**Table 3:** Phytochemicals present in medicinal plants.

Name of medicinal plants	Name of bio-active compounds	Chemical nature of compounds	Analytical technique used	References
<i>Calotropis gigantea</i> (L.) Dryand.	Cardiac glycosides	Calactin, Calotropin, Cymarin, and Asclepin	LC-MS	Bhat and Sharma, 2013
<i>Chamaerops humilis</i> L.	Phenolic compounds	Quinic, Malic acid, Chlorogenic acids, Rutin, and Hesperidin.	LC-MS/MS	Bouhafssoun <i>et al.</i> , 2018
<i>Cuscuta chinensis</i> Lam.	Phenolic acids	Chlorogenic acid, Cryptochlorogenic acid, Neochlorogenic acid, Isochlorogenic acid A, B & C, and Caffeic acid.	HPLC-ESI-MS/MS	Du <i>et al.</i> , 2018
	Flavonoids	Hyperin, Isoquercitrin, Quercetin, Campherol, p-coumaric acid, Isorhamnetin, Rutin, Astragalin, and Apigeni.		
<i>Ephedra alata</i> Decne	Flavonoids	Luteolin-7-O-glucuronide and Myricetin 3-rhamnoside.	HPLC/MS	Al-Rimawi <i>et al.</i> , 2017
<i>Ephedra intermedia</i> Schrenk & C.A.Mey.	Alkaloids	Ephedrine and Pseudoephedrine.	HPLC	Gul <i>et al.</i> , 2017
<i>Ocimum basilicum</i> L.	Terpenoids	$\alpha$ -pinene, $\beta$ -pinene, Limonene, Linalool, and $\alpha$ -terpineol.	GC-MS	Carro <i>et al.</i> , 2013
<i>Phyllanthus emblica</i> L.	Phenolic acid	Gallic acid	LC-MS/MS	Sawant <i>et al.</i> , 2010
	Vitamins	Ascorbic acid		



**Scheme 1:** Phytochemical analysis and antimicrobial activity of bioactive compounds in medicinal plants [Plant extracts (aqueous or alcoholic) are prepared from different fresh or dry parts of the medicinal plant i.e., roots, leaves, barks, flowers, fruits, and seeds. Qualitative and quantitative estimation of the bio-active compounds has been carried out by various bio-chemical tests. Chromatographic methods have been applied to screen the phytochemicals in plant extracts. The antimicrobial activity of plant extracts has been tested using agar dilution method. The MIC/MBC value has been calculated using different concentrations of the extract. The bio-active compounds become fruitful against microbes via inhibiting cell-wall, protein, and nucleic acid synthesis or disrupting the cell-membrane].

It has been observed that leaf extracts of *Acacia nilotica* (L.) Delile, *Sida cordifolia* L., *Tinospora cordifolia* (Willd.) Miers, *W. somnifera* (L.) Dunal and *Ziziphus mauritiana* Lam. exhibited strong antibacterial activity against *Bacillus subtilis*, *Escherichia coli*, *Staphylococcus aureus*, and *Pseudomonas fluorescens* as antifungal activity against *Aspergillus flavus*, *Dreschlera turcica*, and *Fusarium verticilloides* (Mahesh and Satish, 2008). Many polyphenolic constituents and alkaloids are derived from various

medicinal plants, including *Asparagus falcatus* L., *Asteracantha longifolia* Nees, *Vetiveria zizanioides* (L.) Nash, *Epaltes divaricate* (L.) Cass. Moreover, *Coriandrum sativum* L. exhibited excellent antimicrobial activity against *S. aureus* (ATCC 25923), *E. coli* (ATCC 25922) *Pseudomonas aeruginosa* (ATCC 27853), and *Klebsiella pneumoniae* (ATCC 700603) (De Zoysa *et al.*, 2019). The plausible mechanisms of action include cell wall disruption, production of reactive oxygen species, biofilm inhibition, interruption in cell wall construction, DNA replication, energy synthesis, and bacterial toxin synthesis (Mickymaray, 2019). Masoumian and Zandi (2017) observed that hydro-alcoholic extract of *Myrtus communis* L. and aqueous extract of *Cinnamomum zeylanicum* Blume showed strong antibacterial activity against *S. aureus* PTCC 1764, *E. coli* PTCC 1399, *P. aeruginosa* PTCC 1310, and *Salmonella enteric* sub specie PTCC 1709 using disc diffusion method. The extract of *M. communis* L. exhibited the minimum inhibitory concentration (MIC) value at 30 mg/mL compared to penicillin. A study was reported that leaf extract of *M. oleifera* Lam. and flower extract of *Matricaria recutita* L. showed excellent antimicrobial effect against significant drug-resistant clinical isolates such as *P. aeruginosa*, *Staphylococcus* spp., *E. coli*, *Klebsiella* spp., and *Proteus mirabilis* with an MIC value ranging at 7.8–62.5 mg/mL (Atef *et al.*, 2019). Leaf extract of *Combretum collinum* Fresen has been widely used as a traditional medicine to treat wound infection in West Africa. The bio-active compound myricetin-3-O-rhamnoside in the leaf extract of *C. collinum* exhibited most effectiveness against *Staphylococcus epidermidis*, methicillin-resistant *S. aureus* (MRSA), and *S. aureus* (Marquardt *et al.*, 2020). Multi-drug resistance and bacterial biofilm play a significant role in persistent infections, leading to reappearances and deterioration in urinary tract infection (UTI). A study has shown that the essential oil of *Thymus zygis* L., *Origanum majorana* L., and *Rosmarinus officinalis* L. showed antibacterial and anti-biofilm activity against an uropathogen *E. coli* (Lagha *et al.*, 2019). The essential oil of these plants also



exhibited excellent biofilm inhibition and eradication properties against MRSA (Ben Abdallah *et al.*, 2020). Another investigation has shown aqueous acetic acid flower extract of *M. oleifera* Lam. showed strong inhibition effect against multi-drug resistant (MDR) Gram-negative Bacilli (Mandal *et al.*, 2020). Ghosh *et al.* (2014) reported that aqueous leaf extract of *Psidium guajava* L. showed anti-QS activity against an opportunistic pathogen *Chromobacterium violaceum* as manifested by inhibition of violacein production. The leaf extract also inhibits the swarming motility of *P. aeruginosa*. In a review, Gaziano *et al.* (2019) demonstrated that plant extract of *Cardiospermum halicacabum* L. showed excellent antimicrobial activity against both Gram-positive and Gram-negative bacteria. The efficacy of antimicrobial activity may vary plant to plant against organism to organism due to the nature of phytochemicals. A brief account of the antimicrobial activity of different plant extract against various microorganisms is presented in Table 4.

### Antimicrobial Properties of Some Common Medicinal Plants

#### *Allium sativum* L. (Garlic)

*A. sativum* L. has been widely used in Ayurveda due to the presence of allicin. A study has shown that allicin is active against some clinical isolates of lung pathogenic MDR bacteria, including *Pseudomonas*, *Streptococcus*, and *Staphylococcus* (Reiter *et al.*, 2017). Another study suggested that garlic extract play pivotal role in checking the endodontic infections caused by *Porphyromonas gingivalis*, *Aggregatibacter actinomycetemcomitans*, and *Streptococcus mutans* (Hoglund *et al.*, 2020). Nakamoto *et al.* (2020) reported that allicin, vinylidithiols, ajoenes, and diallyl polysulfides are recorded as the

predominant hydrophobic bioactive compounds in garlic and showed promising anti-biofilm activity against Gram-positive and Gram-negative bacteria.

#### *Andrographis paniculata* (Burm.f.) Nees

Ethnic communities have been widely used this plant to cure upper respiratory tract infection and urinary tract infection. A study has shown that the bioactive compound, 14-Deoxy-11, 12-didehydroandrographolide at the concentration of 0.0001M inhibits 92% biofilm production by a 48-hours treatment against *P. aeruginosa* through lessening the content of extracellular polymeric substances, level of pyocyanin production, and synthesis of extracellular protease (Majumdar *et al.*, 2020). Another investigation showed that the plant extract at 100 µg/mL are most effective against CTX-M-15 extended-spectrum β-lactamase-producing *E. coli* (Rasool *et al.*, 2018).

#### *Azadirachta indica* A. Juss (Neem)

The plant has been traditionally used to cure gastrointestinal, uro-genital, and dermatological ailments. The plant extract has noteworthy bactericidal activity against *Helicobacter pylori*. Neem oil extract showed excellent antimicrobial activity against this pathogen with an MIC value between 25–51 µg/mL and MBC value between 43–68 µg/mL (Blum *et al.*, 2019). Mustafa (2016) reported that neem leaf extract showed robust inhibition effect on the growth of *Enterococcus faecalis* compared to 2% chlorhexidine and 3% sodium hypochlorite respectively.

#### *Calotropis gigantea* (L.) Dryand.

The plant has been used as a traditional medicine to treat oral, gastrointestinal, arthritic, and dermal diseases in Indian sub-continent. Aqueous leaf extract of the plant showed excellent

**Table 4:** Antimicrobial activity of plant extract.

Name of the medicinal plants	Use parts	Tested microorganisms	Zone of inhibition (ZOI) (mm)	References
<i>Calotropis gigantea</i> (L.) Dryand.	Milky latex	<i>E. coli</i>	15 ± 1	Rahman <i>et al.</i> , 2013
		<i>Vibrio mimicus</i>	15 ± 1.53	
		<i>Vibrio parahaemolyticus</i>	15 ± 1	
		<i>S. aureus</i>	9 ± 0.58	
<i>Erythrina caffra</i> Thunb.	Bark	<i>Salmonella typhimurium</i>	22 ± 1.3	Wintola <i>et al.</i> , 2021
		<i>P. aeruginosa</i>	35 ± 2.1	
<i>Mentha cervina</i> L.	Aerial parts	<i>S. aureus</i>	30 ± 1.83	Helal <i>et al.</i> , 2019
		<i>Streptococcus pyogenes</i>	19 ± 1.83	
		<i>E. coli</i>	20 ± 0.8	
		<i>S. typhimurium</i>	19 ± 0.82	
<i>Ocimum basilicum</i> L.	Aerial parts	<i>S. aureus</i>	20 ± 2.5	
		<i>Streptococcus pyogenes</i>	19 ± 1.63	
		<i>E. coli</i>	20 ± 1.8	
		<i>S. typhimurium</i>	20 ± 1.83	
<i>Oxalis corniculata</i> L.	Tender shoot	<i>E. coli</i>	17	Manandhar <i>et al.</i> , 2019
		<i>Salmonella typhi</i>	13	
		<i>Salmonella typhi</i> (MDR strain)	16	
		<i>K. pneumoniae</i>	11	
		<i>Citrobacter koseri</i>	12	

antimicrobial activity against various pathogenic strains, including *S. aureus*, *K. pneumoniae*, *Bacillus cereus*, *P. aeruginosa*, *Micrococcus luteus*, and *E. coli* with an MIC value at 50, 25, 6.25, 3.1, 1.5 and 12.5 mg/ml (Kumar *et al.*, 2010). A study has shown that quercetin-3-O-rutinoside has been measured as predominate phytochemical in the plant extract and exhibited remarkable bio-activity against Gram-positive bacteria (*S. aureus* and *B. subtilis*) than the Gram-negative (*P. aeruginosa* and *Salmonella enteritidis*) (Nenaah, 2013).

#### *Catharanthus roseus* (L.) G. Don

The antimicrobial activity of the plant extract was evaluated against pathogenic bacterial strain using agar well diffusion method. The *in-vitro* leaf callus extract exhibited antimicrobial potential against *Bacillus licheniformis* at doses of 2.0 mg/mL (Naz *et al.*, 2015). Khalil (2012) reported that alcoholic extract of plant extract showed magnificent antibacterial activity against *S. aureus* followed by *E. coli* at the dosage of 100 mg/mL. Rani *et al.* (2017) observed that alcoholic method is the best extraction process compared to aqueous, chloroform, and petroleum ether method due to more solubility of bio-active compounds in this plant as because alcoholic extract showed more antimicrobial efficacy than others.

#### *Coccinia grandis* (L.) Voigt

The protease inhibitors (PI), isolated from *Coccinia grandis* (L.) Voigt., have shown antimicrobial action. A study was reported that thermo-stable PI strongly inhibited the growth of pathogenic Gram-positive and Gram-negative microbial strains, including *S. aureus*, *K. pneumoniae*, *Proteus vulgaris*, *E. coli*, and *B. subtilis* (Satheesh and Murugan, 2011). Another investigation showed that aqueous and alcoholic fruit extract showed magnificent antibacterial activity against *S. aureus*, *E. faecalis*, *E. coli*, and *P. aeruginosa* Kirby-Bauer disc diffusion method and compared to erythromycin (Sakharkar and Chauhan, 2017).

#### *Croton bonplandianum* Baill

The plant has been widely used as a traditional medicine for curing of wound infection. The fresh latex of this plant exhibited maximum growth inhibitory activity against enteric bacteria, *E. coli* (ZOI: 32 mm) and *E. faecalis* (ZOI: 30 mm) followed by aqueous and ethanol extracts of latex which exhibited maximum inhibitory activity against *Enterobacter aerogenes* (ZOI: 26 mm) than other solvent extracts. The ethanol and benzene extracts of leaf exhibited antimicrobial activity against *S. aureus* (ZOI: 20 mm). The chloroform extract of fruits showed maximum inhibitory activity against *E. coli* (ZOI: 21 mm) compared to other solvent extracts (Vennila and Udayakumar, 2015).

#### *Curcuma* sp.

Eucalyptol, epicurzerenone, and camphor are recorded as predominant bio-active compounds in the essential oil of *Curcuma caesia* Roxb. The essential oil showed a robust antibacterial activity against *B. subtilis* and *B. cereus* with an MIC at 7.5 µg/mL, while the oil exhibited antifungal activity against *Saccharomyces cerevisiae* with an MIC at 2.5 µg/mL (Paw *et al.*, 2020). *Curcuma longa* L. (turmeric) is enriched with a polyphenolic compound curcumin. It is observed that curcumin caused lipid peroxidation in *S. aureus* and *E. coli*, while the bio-

active compound caused redox homeostasis disparity and triggered the kynurenine pathway in *S. aureus*. Microbial DNA damage occurs due to oxidative stress in exposed curcumin cells (Adeyemi *et al.*, 2020).

#### *Phyllanthus emblica* L. (Amla)

The plant is commonly known as Amla and is used as an important herbal drug in the traditional system of medicine. A report was noted that 1, 2, 3-benzenetriol is screened as a predominant bio-active compound in the fruit extract of amla and showed excellent antimicrobial effect against MDR *Salmonella typhi* and *Salmonella enteritidis* individually or synergistically with *Quercus infectoria* G. Olivier at the doses of 12.5 mg/mL (Nair *et al.*, 2020). A broth dilution study observed that seed extract exhibited quorum-modulatory effect against *Chromobacterium violaceum*, *Serratia marcescens*, *P. aeruginosa*, and *S. aureus* at doses of 50 µg/mL (Patel *et al.*, 2020).

### APPLICATION OF BOTANICAL DISINFECTANT

Plant botanicals are the derivatives of plants or secondary metabolites synthesized by the plant for protective purposes. Botanical disinfectants have been commonly used in various sectors to eliminate pathogens (Hikal *et al.*, 2017). A study was proved that leaf extract of Amla (*P. emblica* L.) checked the growth of *B. subtilis* in mulberry silkworm (*Bombyx mori* L.). The augmentation of silk production is boosted by spraying 10% Amla extract in silk rearing (Gore *et al.*, 2014). Isaiarasu *et al.* (2011) reported that leaf extract of *Acalypha indica* L., *Ocimum sanctum* L. and *Tridax procumbens* (L.) L. controlled the microbial infections in flacherie and muscardine. Apart from mulberry silkworm, plant extract is able to control the flacherie of muga silkworm (*Antheraea assama*). It has been proved that fruit extract of *Terminalia chebula* Retz. (Extract is known as Muga Heal) is very much fruitful against *P. aeruginosa* AC-3, the key pathogen of flacherie in muga silkworm. The gallic acid content in Muga Heal improves the feeding behavior of this worm. It has been found that the accumulation of Muga Heal on the leaves of Som plant (*Machilus bombycina* King ex Hook. f.) increased the quality of cocoons and demolished the rate of pathogenic infection as well (Unni and Neog, 2012). National Research and Development Corporation, India has developed an eco-friendly botanical-based powder (Amruth) to manage grasserie and flacherie diseases. The mixture solution of powder (@2.0g/100 mL) is sprayed uniformly on mulberry leaves (@70.0 mL/kg) before feeding. The leaves are air dried and fed to silkworm (NRDC, 2018). *Bioesque* is a commercial botanical disinfectant and highly effective against a broad spectrum of microorganisms, including bacteria, viruses, and molds. Thymol, a natural monoterpenoid phenol in the oil of thyme, extracted from *Thymus vulgaris* L. is the active ingredient in *Bioesque*. Thymol is low in toxicity to humans, animals, and the environment (Clean Safely, 2020). Therefore, use of botanical disinfectants is a one-step and eco-friendly route to eliminate different pathogens.

### CONCLUSION

Herbal extract has made significant contributions to human health as a therapeutic agent and traditionally offered a source

of opportunity for new drug compounds. Infectious diseases are the leading source of morbidity and mortality across the globe. The numbers of multi-drug resistant bacteria are increasing rapidly due to the indiscriminate use of antibiotics. MDR strains are reduced susceptibility to current antibiotics. The pharmacological use of plant extracts with potential antibacterial activity can be extremely beneficial against microbial pathogens due to noble bio-active compounds. The exploration and conservation of medicinal plants are paramount to explore the investigation on drug designing on this issue. Crude plant extracts have also inhibited the growth of microorganisms through down-regulation of gene expression. It requires more investigation to minimize the emergence of resistance and to establish the most useful therapy for infection. Therefore more plants need to be investigated thoroughly to evaluate the bioactive compound in plant extracts as a potential antimicrobial drug against MDR pathogens.

### Author's Contribution

SS designed the concept. MP and SS contributed equally for writing the final version of the manuscript.

### ACKNOWLEDGMENTS

The authors are grateful to doctors, nurses, paramedical stuff, police personnel, and health workers fighting enormously at the frontline to combat the dreadful SARS- Cov-2.

### REFERENCES

- Q1 Abdelhalim A, Aburjai T, Hanrahan J, Abdel-Halim H. Medicinal plants used by traditional healers in Jordan, the Tafila region. *Pharmacognosy Magazine*. 2017; 13: 95–101. Available from: <https://doi.org/10.4103/0973-1296.203975>
- Adeyemi OS, Obeme-Imom JI, Akpor BO, Rotimi D, Batiha GE, Owolabi A. Altered redox status, DNA damage and modulation of L-tryptophan metabolism contribute to antimicrobial action of curcumin. *Heliyon*. 2020; 6(3): e03495. Available from: <https://doi.org/10.1016/j.heliyon.2020.e03495>
- Ahmad H, Sehgal S, Mishra A, Gupta R. *Mimosa pudica* L. (Laajvanti): An overview. *Pharmacognosy Reviews*. 2012; 6(12): 115–124. Available from: <https://dx.doi.org/10.4103%2F0973-7847.99945>
- Ahmed E, Arshad M, Khan MZ, Amjad MS, Sadaf HM, Riaz I, Sabir S, Ahmad N. Secondary metabolites and their multidimensional prospective in plant life. *Journal of Pharmacognosy and Phytochemistry*. 2017; 6: 205–214.
- Al-Rimawi F, Abu-Lafi S, Abbadi J, Alamarneh AAA, Sawahreh RA, Odeh I. Analysis of phenolic and flavonoids of wild *Ephedra alata* plant extract by LC/PDA and LC/MS and their antioxidant activity. *African Journal of Traditional, Complementary, and Alternative Medicines*. 2017; 14: 130–141. Available from: <https://dx.doi.org/10.21010%2Fajtcam.v14i2.14>
- Atef NM, Shanab SM, Negm SI, Abbas YA. Evaluation of antimicrobial activity of some plant extracts against antibiotic susceptible and resistant bacterial strains causing wound infection. *Bulletin of the National Research Centre*. 2019; 43: 144. Available from: <https://doi.org/10.1186/s42269-019-0184-9>
- Aziz MA, Adnan M, Khan AH, Shahat AA, Al-Said MS, Ullah R. Traditional uses of medicinal plants practiced by the indigenous communities at Mohmand Agency, FATA, Pakistan. *Journal of Ethnobiology and Ethnomedicine*. 2018; 14: 2. Available from: <https://doi.org/10.1186/s13002-017-0204-5>
- Bacha K, Tariku Y, Gebreyesus F, Zerihun S, Mohammed A, Weiland-Bräuer N, Schmitz RA, Mulat M. Antimicrobial and anti-Quorum Sensing activities of selected medicinal plants of Ethiopia: Implication for development of potent antimicrobial agents. *BMC Microbiology*. 2016; 16(1): 139. Available from: <https://doi.org/10.1186/s12866-016-0765-9>
- Ben Abdallah, F, Lagha R, Gaber A. 2020. Biofilm Inhibition and Eradication Properties of Medicinal Plant Essential Oils against Methicillin-Resistant *Staphylococcus aureus* Clinical Isolates. *Pharmaceuticals* (Basel, Switzerland). 13(11): 369. Available from: <https://doi.org/10.3390/ph13110369>
- Bhat KS, Sharma A. Therapeutic potential of cardiac glycosides of *Calotropis gigantea* for breast cancer. *International Research Journal of Pharmacy*. 2013; 4: 164–167. Available from: <http://dx.doi.org/10.7897/2230-8407.04636>
- Bhavana KR, Shreevathsa. Medical geography in Charaka Samhita. *Ayu*. 2014; 35(4): 371–377. Available from: <http://dx.doi.org/10.4103/0974-8520.158984>
- Blum FC, Singh J, Merrell DS. In vitro activity of neem (*Azadirachta indica*) oil extract against *Helicobacter pylori*. *Journal of Ethnopharmacology*. 2019; 232: 236–243. Available from: <https://doi.org/10.1016/j.jep.2018.12.025>
- Bouhafssoun A, Yilmaz MA, Boukelou A, Temel H, Harche MK. Simultaneous quantification of phenolic acids and flavonoids in *Chamaerops humilis* L. using LC–ESI-MS/MS. *Food Science and Technology*. 2018; 38: 242–247. Available from: <http://dx.doi.org/10.1590/fst.19917>
- Carro MD, Ianni C, Magi E. Determination of terpenoids in plant leaves by GC-MS: development of the method and application to *Ocimum basilicum* and *Nicotiana glauca*. *Analytical Letters*. 2013; 46: 630–639. Available from: <https://doi.org/10.1080/00032719.2012.729239>
- Clean S. Bioesque Disinfectant Solution Overview. 2020. Available from: <https://www.clean-safely.com/bioesque-disinfectant-solution-overview/>
- Das K, Tiwari RKS, Shrivastava DK. Techniques for evaluation of medicinal plant products as antimicrobial agents: current methods and future trends. *Journal of Medicinal Plants Research*. 2010; 4: 104–111. Available from: <http://www.academicjournals.org/JMPR>
- De Zoysa MHN, Rathnayake H, Hewawasam RP, Gaya Bandara Wijayarathne WMD. Determination of *in-vitro* antimicrobial activity of five Sri Lankan medicinal plants against selected human pathogenic bacteria. *International Journal of Microbiology*. 2019; 2019: 7431439. Available from: <https://doi.org/10.1155/2019/7431439>
- Dhandapani R, Sabna B. Phytochemical constituents of some Indian medicinal plants. *Ancient Science of Life*. 2008; 27: 1–8.
- Dolatkhahi M, Dolatkhahi A, Nejad JB. Ethnobotanical study of medicinal plants used in Arjan - Parishan protected area in Fars Province of Iran. *Avicenna Journal of Phytomedicine*. 2014; 4(6): 402–412.
- Du K, Li J, Guo X, Li Y, Chang Y. Quantitative analysis of phenolic acids and flavonoids in *Cuscuta chinensis* Lam. by synchronous ultrasonic-assisted extraction with response surface methodology. *Journal of Analytical Methods in Chemistry*. 2018; 2018: 6796720. Available from: <http://dx.doi.org/10.1155/2018/6796720>
- Elisha IL, Botha FS, McGaw LJ, Eloff JN. The antibacterial activity of extracts of nine plant species with good activity against *Escherichia coli* against five other bacteria and cytotoxicity of extracts. *BMC Complementary and Alternative Medicine*. 2017; 17(1): 133. Available from: <http://doi.org/10.1186/s12906-017-1645-z>
- Fahal EM, Babitha Rani AM, Aklakur MD, Chanu TI, Saharan N. Qualitative and quantitative phytochemical analysis of *Moringa oleifera* (Lam) pods. *International Journal of Current Microbiology and Applied Sciences*. 2018; 7: 657–665. Available from: <http://dx.doi.org/10.20546/ijcmas.2018.705.080>
- Fraga CG, Croft KD, Kennedy DO, Tomás-Barberán FA. The effects of polyphenols and other bioactives on human health. *Food & Function*. 2019; 10(2): 514–528. Available from: <http://doi.org/10.1039/c8fo01997e>
- Gandhiraja N, Sriram S, Meenaa V, Srilakshmi JK, Sasikumar C, Rajeswari R. Phytochemical Screening and Antimicrobial Activity of the Plant Extracts of *Mimosa pudica* L. Against Selected Microbes. *Ethnobotanical Leaflets*. 2009; 5: 8.
- Gaziano R, Campione E, Iacovelli F, Pistoia ES, Marino D, Milani M, Di Francesco P, Pica F, Bianchi L, Orlandi A, Marsico S, Falconi M, Aquaro



- S. Antimicrobial properties of the medicinal plant *Cardiospermum halicacabum* L: new evidence and future perspectives. *European Review for Medical and Pharmacological Sciences*. 2019; 23(16): 7135–7143. Available from: [http://doi.org/10.26355/eurrev\\_201908\\_18759](http://doi.org/10.26355/eurrev_201908_18759)
- Ghosh R, Tiwary BK, Kumar A, Chakraborty R. Guava Leaf Extract Inhibits Quorum-Sensing and *Chromobacterium violaceum* Induced Lysis of Human Hepatoma Cells: Whole Transcriptome Analysis Reveals Differential Gene Expression. *PLoS ONE*. 2014; 9(9): e107703. Available from: <https://doi.org/10.1371/journal.pone.0107703>
- Gore D, Rai M, Shinde G. Effect of Amla extract on bacterial infection in the silkworm, *Bombyx mori* L. *Journal of Pharmacy Research*. 2014; 8(5): 679–680. Available from: <http://jprsolution.info/>
- Gul R, Jan SU, Faridullah S, Sherani S, Jahan N. Preliminary phytochemical screening, quantitative analysis of alkaloids, and antioxidant activity of crude plant extracts from *Ephedra intermedia* indigenous to Balochistan. *The Scientific World Journal*. 2017; 2017: 5873648. Available from: <https://doi.org/10.1155/2017/5873648>
- Helal IM, El-Bessoumy A, Al-Bataineh E, Joseph M, Rajagopalan P, Chandramoorthy HC, Ben Hadj AS. Antimicrobial Efficiency of Essential Oils from Traditional Medicinal Plants of Asir Region, Saudi Arabia, over Drug Resistant Isolates. *BioMed Research International*. 2019; 2019: 8928306. Available from: <https://doi.org/10.1155/2019/8928306>
- Hikal WM, Baeshen RS, Said-Al Ahl HAH. Botanical insecticides as simple extractives for pest control. *Cogent Biology*. 2017; 3(1): 1404274. Available from: <https://doi.org/10.1080/23312025.2017.1404274>
- Hoglund KB, Barnett BK, Watson SA, Melgarejo MB, Kang Y. Activity of bioactive garlic compounds on the oral microbiome: a literature review. *General Dentistry*. 2020; 68(3): 27–33.
- Isaiarasu L, Sakthivel N, Ravikumar J, Samuthiravelu P. Effect of herbal extracts on the microbial pathogens causing flacherie and muscardine diseases in the mulberry silkworm, *Bombyx mori* L. *Journal of Biopesticides*. 2011; 4(2): 150–155.
- Ismandari T, Kumalaningsih S, Wijana S, Mustanirah SA. Optimization of bioactive compound extraction from rose myrtle fruit (*Rhodomyrtus tomentosa*, (W.Ait), *Myrtaceae*) as the antioxidant source. *The Scientific World Journal*. 2020; 2020: 9105847. Available from: <https://doi.org/10.1155/2020/9105847>
- Jindal A, Kumar P, Singh G. Extraction and pharmacological evaluation of flavonoids of *Sida acuta* Burm. F. *International journal of Green Pharmacy*. 2012; 6: 208–211. Available from: <http://dx.doi.org/10.22377/ijgp.v6i3.263>
- Jolly CI, Narayanan P. Pharmacognosy of aerial parts of *Cynodon dactylon* Pers. (Graminae). *Ancient Science of Life*. 2000; 19: 1–6.
- Khalil A. Antimicrobial activity of ethanol leaf extracts of *Catharanthus roseus* from Saudi Arabia. Singapore: Second International Conference on Environment Science and Biotechnology. 2012; 48.
- Kumar G, Karthik L, Rao KVB. Antibacterial activity of aqueous extract of *Calotropis gigantea* leaves – an in-vitro study. *International Journal of Pharmaceutical Sciences Review and Research*. 2010; 4(2): 141–144.
- Lagha R, Ben Abdallah F, Al-Sarhan BO, Al-Sodany Y. Antibacterial and Biofilm Inhibitory Activity of Medicinal Plant Essential Oils against *Escherichia coli* Isolated from UTI Patients. *Molecules (Basel, Switzerland)*. 2019; 24(6): 1161. Available from: <https://doi.org/10.3390/molecules24061161>
- Mahasneh AM, El-Oqlah AA. Antimicrobial activity of extracts of herbal plants used in the traditional medicine of Jordan. *Journal of Ethnopharmacology*. 1999; 64(3): 271–276. Available from: [https://doi.org/10.1016/S0378-8741\(98\)00132-9](https://doi.org/10.1016/S0378-8741(98)00132-9)
- Mahesh B, Satish S. Antimicrobial activity of some important medicinal plant against plant and human pathogens. *World Journal of Agricultural Sciences*. 2008; 4: 839–843.
- Mahmoud T, Gairola S. Traditional knowledge and use of medicinal plants in the Eastern desert of Egypt: a case study from Wadi El-Gemal National Park. *Journal of Medicinal Plants Studies*. 2013; 1: 10–17.
- Mahomoodally MF. Traditional medicines in Africa: an appraisal of ten potent African medicinal plants. *Evidence-Based Complementary and Alternative Medicine*. 2013; 2013: 617459. Available from: <https://doi.org/10.1155/2013/617459>
- Mahwasane ST, Middleton L, Boaduo N. An ethnobotanical survey of indigenous knowledge on medicinal plants used by the traditional healers of the Lwamondo area, Limpopo province, South Africa. *South African Journal of Botany*. 2013; 88: 69–75. Available from: <https://doi.org/10.1016/j.sajb.2013.05.004>
- Mailu JK, Nguta JM, Mbaria JM, Okumu MO. Medicinal plants used in managing diseases of the respiratory system among the Luo community: an appraisal of Kisumu East Sub-County, Kenya. *Chinese Medicine*. 2020; 15: 95. Available from: <https://doi.org/10.1186/s13020-020-00374-2>
- Majumdar M, Misra TK, Roy DN. In-vitro anti-biofilm activity of 14-deoxy-11, 12-didehydroandrographolide from *Andrographis paniculata* against *Pseudomonas aeruginosa*. *Brazilian Journal of Microbiology*: [publication of the Brazilian Society for Microbiology]. 2020; 51(1): 15–27. Available from: <https://dx.doi.org/10.1007/s2020-019-00169-0>
- Manandhar S, Luitel S, Dahal RK. In Vitro Antimicrobial Activity of Some Medicinal Plants against Human Pathogenic Bacteria. *Journal of Tropical Medicine*. 2019; 2019: 1895340. <https://doi.org/10.1155/2019/1895340>
- Mandal SM, Chakraborty S, Sahoo S, Pyne S, Ghosh S, Chakraborty R. Novel Compound from Flowers of *Moringa oleifera* Active against Multi-Drug Resistant Gram-negative Bacilli. *Infectious Disorders Drug Targets*. 2020; 20(1): 69–75. Available from: <https://doi.org/10.2174/1871526518666181001124420>
- Marquardt P, Seide R, Vissienon C, Schubert A, Birkemeyer C, Ahyi V, Fester K. Phytochemical Characterization and In Vitro Anti-Inflammatory, Antioxidant and Antimicrobial Activity of *Combretum collinum* Fresen Leaves Extracts from Benin. *Molecules (Basel, Switzerland)*. 2020; 25(2): 288. Available from: <https://doi.org/10.3390/molecules25020288>
- Masoumian M, Zandi M. Antimicrobial Activity of Some Medicinal Plant Extracts against Multidrug Resistant Bacteria. *Zahedan Journal of Research in Medical Sciences*. 2017; 19(11): e10080. Available from: <https://dx.doi.org/10.5812/zjrms.10080>
- Mickymaray S. Efficacy and mechanism of traditional medicinal plants and bioactive compounds against clinically important pathogens. *Antibiotics (Basel, Switzerland)*. 2019; 8: 257. Available from: <https://doi.org/10.3390/antibiotics8040257>
- Mishra R, Pounikar Y, Gangwar M. Extraction, Phytochemical screening and quantitative determination of phenols and flavonoids in extract of *Kalanchoe pinnata* and *Pongamia pinnata*. *Journal of Drug Delivery and Therapeutics*. 2019; 9: 192–196. Available from: <https://doi.org/10.22270/jddt.v9i4.3024>
- Mishra S, Aeri V, Gaur PK, Jachak SM. Phytochemical, therapeutic, and ethnopharmacological overview for a traditionally important herb: *Boerhavia diffusa* Linn. *BioMed Research International*. 2014; 2014: 808302. Available from: <https://doi.org/10.1155/2014/808302>
- Mngeni NZ. Bio-active compounds from selected medicinal plants used in anti-diabetic treatment. M. Tech. Thesis in Chemistry, Cape Peninsula University of Technology. 2017.
- Morales F, Padilla S, Falconí F. Medicinal plants used in traditional herbal medicine in the province of Chimborazo, Ecuador. *African Journal of Traditional, Complementary, and Alternative Medicines*. 2016; 14: 10–15. Available from: <https://dx.doi.org/10.21010/2Fajtcam.v14i1.2>
- Mustafa M. Antibacterial Efficacy of Neem (*Azadirachta indica*) Extract against *Enterococcus faecalis*: An in vitro Study. *The Journal of Contemporary Dental Practice*. 2016; 17(10): 791–794. Available from: <https://doi.org/10.5005/jp-journals-10024-1932>
- Nair A, Balasaravanan T, Jadhav S, Mohan VKC. Harnessing the antibacterial activity of *Quercus infectoria* and *Phyllanthus emblica* against antibiotic-resistant *Salmonella Typhi* and *Salmonella Enteritidis* of poultry origin. *Veterinary World*. 2020; 13(7): 1388–1396. Available from: <https://doi.org/10.14202/vetworld.2020.1388-1396>
- Nakamoto M, Kunitura K, Suzuki JI, Kodera Y. Antimicrobial properties of hydrophobic compounds in garlic: Allicin, vinylthiins, ajoene and diallyl polysulfides. *Experimental and Therapeutic Medicine*. 2020; 19(2): 1550–1553. Available from: <https://dx.doi.org/10.3892/etm.2019.8388>



- Nandagoapalan V, Marimuthuand C, Doss A. Diversity of traditional medicinal plants used by rural community in Tiruchirappalli district Tamilnadu, South India. *International Journal of Current Microbiology and Applied Sciences*. 2015; 4: 767-776.
- National Research and Development Corporation (NRDC), India. Amruth–A botanical product for controlling grasserie and flacherie in silkworm. 2018. Available from: <https://www.nrdcindia.com/english/index.php/...05.../sericulture/305-seri-tech-7>
- Nawrot R, Barylski J, Nowicki G, Broniarczyk J, Buchwal W, Goździcka-Józefiak A. Plant antimicrobial peptides. *Folia Microbiologica*. 2014; 59(3): 181–196. Available from: <https://doi.org/10.1007/s12223-013-0280-4>
- Naz S, Haq R, Aslam F, Ilyas S. Evaluation of antimicrobial activity of extracts of *in-vivo* and *in vitro* grown *Vinca rosea* L. (*Catharanthus roseus*) against pathogens. *Pakistan Journal of Pharmaceutical Sciences*. 2015; 28(3): 849–853.
- Neba NE. Traditional Health Care System and Challenges in Developing Ethnopharmacology in Africa: Example of Oku, Cameroon. *Studies on Ethno-Medicine*. 2011; 5(2): 133-139. Available from: <https://doi.org/10.1080/09735070.2011.11886400>
- Nenaah G. Antimicrobial activity of *Calotropis procera* Ait. (Asclepiadaceae) and isolation of four flavonoid glycosides as the active constituents. *World Journal of Microbiology & Biotechnology*. 2013; 29(7): 1255–1262. Available from: <https://doi.org/10.1007/s11274-013-1288-2>
- Oladeji O. The Characteristics and Roles of Medicinal Plants: Some Important Medicinal Plants in Nigeria. *Natural Products: An Indian Journal*. 2016; 12(3): 10.
- Osman A, Sbhatu DB, Giday M. Medicinal Plants Used to Manage Human and Livestock Ailments in Raya Kobo District of Amhara Regional State, Ethiopia. *Evidence-Based Complementary and Alternative Medicine: eCAM*. 2020; 2020: 1329170. Available from: <https://doi.org/10.1155/2020/1329170>
- Pan SY, Litscher G, Gao SH, Zhou SF, Yu ZL, Chen HQ, Zhang SF, Tang MK, Sun JN, Ko KM. Historical perspective of traditional indigenous medical practices: the current renaissance and conservation of herbal resources. *Evidence-Based Complementary and Alternative Medicine: eCAM*. 2014; 2014: 525340. Available from: <https://doi.org/10.1155/2014/525340>
- Pandey MM, Rastogi S, Rawat AKS. Indian Traditional Ayurvedic System of Medicine and Nutritional Supplementation. *Evidence-Based Complementary and Alternative Medicine*. 2013; 2013: 376327. Available from: <https://doi.org/10.1155/2013/376327>
- Patel P, Joshi C, Kothari V. The Anti-infective Potential of Hydroalcoholic Extract of *Phyllanthus emblica* Seeds Against Selected Human-pathogenic Bacteria. *Infectious Disorders Drug Targets*. 2020; 20(5): 672–692. Available from: <https://doi.org/10.2174/1871526519666190821154926>
- Paw M, Gogoi R, Sarma N, Pandey SK, Borah A, Begum T, Lal M. Study of Anti-oxidant, Anti-inflammatory, Genotoxicity, and Antimicrobial Activities and Analysis of Different Constituents found in Rhizome Essential Oil of *Curcuma caesia* Roxb., Collected from North East India. *Current Pharmaceutical Biotechnology*. 2020; 21(5): 403–413. <https://doi.org/10.2174/1389201020666191118121609>
- Petrovska BB. Historical review of medicinal plants' usage. *Pharmacognosy Reviews*. 2012; 6(11): 1–5. Available from: <https://doi.org/10.4103/0973-7847.95849>
- Poonia A. Bioactive Compounds of Fat-Hen (*Chenopodium album* L.). In: Murthy HN, Paek KY, editors. *Bioactive Compounds in Underutilized Vegetables and Legumes*. Switzerland: Springer Nature AG; 2021, 1-11 pp.
- Prasad LV. In: Chaudhury RR, Rafei UM editors. *Indian System of Medicine and Homoeopathy Traditional Medicine in Asia*. New Delhi (India): WHO- Regional Office for South East Asia; 2002, 283–286 pp.
- Rahman MS, Moly NN, Hossen MJ. Antimicrobial, Cytotoxic and Antioxidant activity of the exudate of *Calotropis gigantea*. *International Journal of Pharmaceutical Sciences and Research*. 2013; 4(2): 745-753.
- Rajan S, Gokila A, Jency P, Brindha P, Sujatha, RK. Antioxidant and phytochemical properties of *Aegle marmelos* fruit pulp. *International Journal of Current Pharmaceutical Research*. 2011; 3: 65–70.
- Rajendran R, Sundararajan R. Preliminary Phytochemical Analysis and Anti-Bacterial Activity of *Mimosa pudica* Linn. Leaves. *International Journal of Pharma and Bio Sciences*. 2010; 6: 1–8.
- Rani J, Kapoor M, Kaur R. In-vitro anti-bacterial activity and phytochemical screening of crude extracts of *Catharanthus roseus* L. (G.) Don. *Agricultural Science Digest*. 2017; 37(2): 106-111.
- Rasool U, S P, Parveen A, Sah SK, S H. Efficacy of *Andrographis paniculata* against extended spectrum  $\beta$ -lactamase (ESBL) producing *E. coli*. *BMC Complementary and Alternative Medicine*. 2018; 18(1): 244. Available from: <https://doi.org/10.1186/s12906-018-2312-8>
- Ravishankar B, Shukla VJ. Indian systems of medicine: a brief profile. *African Journal of Traditional, Complementary, and Alternative Medicines: AJTCAM*. 2007; 4(3): 319–337. Available from: <https://doi.org/10.4314/ajtcam.v4i3.31226>
- Reiter J, Levina N, van der Linden M, Gruhlke M, Martin C, Slusarenko AJ. Diallylthiosulfinate (Allicin), a Volatile Antimicrobial from Garlic (*Allium sativum*), Kills Human Lung Pathogenic Bacteria, Including MDR Strains, as a Vapor. *Molecules* (Basel, Switzerland). 2017; 22(10): 1711. Available from: <https://doi.org/10.3390/molecules22101711>
- Saini S, Dhiman A, Nanda S. Traditional Indian medicinal plants with potential wound healing activity: a review. *International Journal of Pharmaceutical Sciences and Research*. 2016; 7: 1809-1819.
- Sakharkar P, Chauhan B. Antibacterial, antioxidant and cell proliferative properties of *Coccinia grandis* fruits. *Avicenna Journal of Phytomedicine*. 2017; 7(4): 295–307.
- Satheesh LS, Murugan K. Antimicrobial activity of protease inhibitor from leaves of *Coccinia grandis* (L.) Voigt. *Indian Journal of Experimental Biology*. 2011; 49(5): 366–374.
- Sawant L, Prabhakar B, Pandita N. Quantitative HPLC analysis of ascorbic acid and gallic acid in *Phyllanthus emblica*. *Journal of Analytical and Bio-analytical Techniques*. 2010; 1: 111.
- Shaikh SS. Preliminary phytochemical qualitative and quantitative screening of *Tridax procumbens* L. *International Journal of Chemical and Physical Sciences*. 2018; 7: 213-217.
- Some S, Mukherjee J. Study on Some Leafy Vegetables and their Medicinal Uses at Chanchal Sub - Division of Malda District, West Bengal. *International Journal of Plant and Environment*. 2018; 4(1): 97-104. Available from: <https://ijplantenviro.com/index.php/IJPE/article/view/1044>
- Some S, Ghosh A, Mukherjee J, Basu D. Study of some traditionally important medicinal plants for primary healthcare by local people in the northern part of Malda district, West Bengal (India). *International Journal of Current Research and Academic Review*. 2017; 5(7):1-9. Available from: <https://doi.org/10.20546/ijcrar.2017.507.001>
- Srinivasan D, Perumalsamy LP, Nathan S, Sures T. Antimicrobial Activity of Certain Indian Medicinal Plants Used in Folkloric Medicine. *Journal of Ethnopharmacology*. 2001; 74(3): 217-222. Available from: [https://doi.org/10.1016/s0378-8741\(00\)00345-7](https://doi.org/10.1016/s0378-8741(00)00345-7)
- Tefera BN, Kim YD. Ethnobotanical study of medicinal plants in the Hawassa Zuria District, Sidama zone, Southern Ethiopia. *Journal of Ethnobiology and Ethnomedicine*. 2019; 15(1): 25. Available from: <https://doi.org/10.1186/s13002-019-0302-7>
- Teka A, Asfaw Z, Demissew S, Van Damme P. Medicinal plant use practice in four ethnic communities (Gurage, Mareqo, Qebena, and Silti), south central Ethiopia. *Journal of Ethnobiology and Ethnomedicine* 2020; 16(1): 27. Available from: <https://doi.org/10.1186/s13002-020-00377-1>
- Thirumalai T, Kelumalai E, Senthilkumar B, David E. Ethnobotanical study of medicinal plants used by the local people in Vellore district, Tamilnadu, India. *Ethnobotanical Leaflets*. 2009; 2009(10): 10. Available from: <https://opensiuc.lib.siu.edu/ebf/vol2009/iss10/10>
- Uniyal SK, Singh KN, Jamwal P, Lal, B. Traditional use of medicinal plants among the tribal communities of Chhota Bhangal, Western Himalaya. *Journal of Ethnobiology and Ethnomedicine*. 2006; 2: 14. Available from: <https://doi.org/10.1186/1746-4269-2-14>
- Unni BG, Neog K. Current trend of biotechnological and molecular research in Muga silk worm, *Antheraea assamensis*, helper and future prospects. Guwahati, Assam, India: National Seminar on “Recent Trends in Research & Development in Muga Culture – Ideas to Action”; 2012, 12-20 pp.

- Upreti Y, Asselin H, Dhakal A, Julien N. Traditional use of medicinal plants in the boreal forest of Canada: review and perspectives. *Journal of Ethnobiology and Ethnomedicine*. 2012; 8: 7. Available from: <https://doi.org/10.1186/1746-4269-8-7>
- Vennila V, Udayakumar R. Antibacterial activity of *Croton bonplandianum* (Bail.) against some bacterial isolates from infected wounds. *British Microbiology Research Journal* 2015; 5: 83-93. Available from: <http://doi.org/10.9734/BMRJ/2015/12620>
- Vidyarthi S, Samant SS, Sharma P. Traditional and indigenous uses of medicinal plants by local residents in Himachal Pradesh, North Western Himalaya, India. *International Journal of Biodiversity Science, Ecosystem Services & Management*. 2013; 9: 185-200. Available from: <https://doi.org/10.1080/21513732.2013.823113>
- White P. Integron and Antibiotic Resistance. *Pathology*. 2009; 41(S1): 46.
- Wintola, OA, Olajuyigbe, AA, Afolayan AJ, Cooposamy RM, Olajuyigbe OO. Chemical composition, antioxidant activities and antibacterial activities of essential oil from *Erythrina caffra* Thunb. growing in South Africa. *Heliyon*. 2021; 7(6): e07244. Available from: <https://doi.org/10.1016/j.heliyon.2021.e07244>
- Yadav NK, Arya RK, Dev K, Sharma C, Hossain Z, Meena S, Singh RK. Alcoholic extract of *Eclipta alba* shows *in-vitro* antioxidant and anticancer activity without exhibiting toxicological effects. *Oxidative Medicine and Cellular Longevity*. 2017; 2017: 9094641. Available from: <https://doi.org/10.1155/2017/9094641>
- Yahia Y, Benabderrahim MA, Tlili N, Bagues M, Nagaz K. Bioactive compounds, antioxidant and antimicrobial activities of extracts from different plant parts of two *Ziziphus* Mill. Species. *PLoS One*. 2020; 15: e0232599. Available from: <https://doi.org/10.1371/journal.pone.0232599>

#### Query Report

Q1. Cross-check for accuracy/data/format, correct, and provide all references in consistent Vancouver format.  
 Sample reference formatted to act as an example and to maintain consistency of format (Vancouver) required  
 Kumar RA, Sridevi K, Kumar NV, Nanduri S, Rajagopal S. Anticancer and immunostimulatory compounds from *Andrographis paniculata*. *Journal of Ethnopharmacology*. 2004;92:291-295. Available from: [doi.org/10.1016/j.jep.2004.03.004](https://doi.org/10.1016/j.jep.2004.03.004)